

Production of photons in a bouncing universe

J. M. Salim¹, S. E. Perez Bergliaffa^{1,2} and N. Souza²

¹*Centro Brasileiro de Pesquisas Físicas, Rua Xavier Sigaud,
150, CEP 22290-180, Rio de Janeiro, Brazil, and*

²*Departamento de Física Teórica, Instituto de Física,
Universidade do Estado de Rio de Janeiro, CEP 20550-013, Rio de Janeiro, Brazil*

Using a new non-singular solution, it is shown that the production of photons in dilaton electrodynamics in a cosmological setting is increased if the effect of matter creation on the geometry is taken into account. This increment may be related to the problem of the origin of magnetic fields in the universe.

Introduction

The origin, evolution, and structure of the magnetic fields present in galaxies and clusters of galaxies are amongst the most important open issues in astrophysics and cosmology. Typically, the magnetic field present in galaxies is of the order of a few μG , and is coherent on the galactic scale. The observed field in the case of clusters is correlated over 10-100 kiloparsecs, and can be as high as tens of μG . The standard mechanism to account for these fields is the dynamo [1], which amplifies small seed fields to the abovementioned values [2]. The seeds that fuel the dynamo may have an astrophysical origin, or may be primordial. An astrophysical mechanism for the generation of these pre-galactic magnetic fields is the Biermann battery [3], which works for the generation of seeds to be amplified by the dynamo in galaxies, but can hardly account for the fields in clusters [4, 5]. Other astrophysical mechanisms (involving for instance starbursts, jet-lobe radio sources, or accretion disks of black holes in AGNs [6]) seem to need pre-existing magnetic fields. The current prevalent view is that the magnetic fields observed in galaxies and clusters have a primordial origin (see [7] for a review). The processes that may account for this origin can be divided in two types: causal (those in which the seeds are produced at a given time inside the horizon, like QCD and EW phase transitions [7]), and inflationary (where correlations are produced outside the horizon [8]). In the latter, vacuum fluctuations of the electromagnetic field are “stretched” by the evolution of the background geometry to super-horizon scales, and they could appear today as large-scale magnetic fields. However, since Maxwell’s equations are conformally invariant in the FRW background, the amplification of the vacuum fluctuations (which amounts to particle production) via inflation can work only if conformal invariance is broken at some stage of the evolution of the universe. There are several ways in which this invariance can be broken [9]: non-minimal coupling between gravitation and the electromagnetic field [8, 10, 11], quantum anomaly of the trace of the stress-energy tensor of electrodynamics [12], coupling of the EM field to a charged scalar field [8, 13], exponential coupling between a scalar (whose potential

drives inflation) and the EM field [14], and a non-zero mass for the photon [15]. Yet another possibility is dilaton electrodynamics [16], in which there is a scalar field (the dilaton) which couples exponentially to an Abelian gauge field. In this model the inflationary expansion is driven and not by the potential but by the kinetic term of the scalar field. The exponential coupling is naturally implemented in the low-energy limit in string theory [17], and in Weyl integrable spacetime (WIST) [18], and can be viewed also as a time dependence of the coupling constant, an idea considered first by Dirac [19]. This avenue has been pursued by Giovannini in a series of articles [20]. More recently, a model with both a dilaton and an inflaton was analyzed in [21], and later generalized to a noncommutative spacetime [22]. In all these articles, different aspects of photon production have been analyzed. We will focus in this contribution on two definite items. First, we shall address the creation of photons in a non-singular universe described by an exact solution of the field equations. More specifically, in this solution the passage through the bounce is described by the equations of the model without resorting to unknown (Planck-scale) physics. The results of this calculation are to be compared with those of the second item, coming from a more complete model. We shall take into account in this second part the influence of the creation of matter in the squeezing of the vacuum state, a problem that seems to have received little attention in the literature. It will be shown that as a result of matter creation, there is an increment in the number of photons originating in the stretching of the vacuum fluctuations. Thus this effect may be of importance in the generation of primordial magnetic fields. The analysis will be done in the framework of a phenomenological model, in which the creation of ultrarelativistic matter is restricted to an interval centered in the bounce. Two interesting features of the solution representing this model are that it goes automatically into a radiation regime after a short time, and it displays a constant value for the dilaton after entering the radiation era. We shall start in Sect.1 by deducing a Hamiltonian for the EM field in a curved background coupled to the scalar field. The quantization of this Hamiltonian system will be presented in Sect.2, following the method of the squeezed states. In Sect.3 we

shall study the production of photons in the absence of matter (*i.e.* with only the scalar field present). Variation of the photon number in a model that takes into account matter creation will be discussed in Sect.4. We close with a discussion in Sect.5.

I. FIELD EQUATIONS

A time-dependent gauge coupling is a generic feature of 4-dimensional theories obtained by compactification of some theory formulated in a spacetime with more than 4 dimensions, such as Kaluza-Klein theory [23] and string theory [24]. Time-dependent gauge couplings are also present in WIST [18]. In all these cases, the action can be conveniently written in the form

$$S = \frac{1}{2} \int d^4x \sqrt{-g} f(\omega) F_{\alpha\beta} F^{\alpha\beta}, \quad (1)$$

where ω is the the dilaton (in the case of string theory) or the scalar field associated to Weyl geometry (in the case of WIST) and $F_{\mu\nu}$ is an Abelian field. The function $f(\omega)$ will be left unspecified for the time being, but later we will set $f(\omega) = e^{-2\omega}$, which corresponds to the case of string theory and WIST. The equations of motion (EOM) that follow from this action are:

$$(f(\omega) F^{\mu\nu})_{||\nu} = 0, \quad (2)$$

$$F^{*\alpha\beta}_{||\beta} = 0, \quad (3)$$

where the derivative wrt the background geometry is denoted by " $||$ ", and the dual of the electromagnetic tensor $F_{\mu\nu}$ is defined as

$$F^{*\alpha\beta} \equiv \frac{1}{2} \eta^{\alpha\beta\mu\nu} F_{\mu\nu}.$$

Advantage will be taken in the following of the formal equivalence of Eqns.(2) and (3) and the equations for the electromagnetic field in a material medium, an equivalence that is fulfilled if we define

$$P^{\mu\nu} \doteq f(\omega) F^{\mu\nu}. \quad (4)$$

A reference frame must be chosen in order to define the electric and magnetic fields from $P_{\mu\nu}$ and $F_{\mu\nu}$. Since the cosmological model to be used in this article is described by the Friedmann-Robertson-Walker (FRW) metric, we can use the four velocity vector field $V^\mu = \delta^\mu_0$, which is orthogonal to the three-dimensional surfaces of homogeneity of the FRW geometry. The electric and magnetic parts of the electromagnetic tensor are defined as:

$$D^\alpha \doteq P^{\alpha\beta} V_\beta, \quad H^\alpha \doteq P^{*\alpha\beta} V_\beta, \quad (5)$$

$$E^\alpha \doteq F^{\alpha\beta} V_\beta, \quad B^\alpha \doteq F^{*\alpha\beta} V_\beta. \quad (6)$$

In the comoving frame, the EOM along with Eqn.(4) become

$$D^\alpha_{||\alpha} = 0, \quad (7)$$

$$B^\alpha_{||\alpha} = 0, \quad (8)$$

$$(h^\alpha_\lambda D^\alpha) + \frac{2}{3} \theta D_\lambda + \eta_\lambda^{\beta\rho\sigma} V_\rho H_{\sigma||\beta} = 0, \quad (9)$$

$$h^\alpha_\lambda \dot{B}^\alpha + \frac{2}{3} \theta B_\lambda + \eta_\lambda^{\beta\rho\sigma} V_\rho E_{\sigma||\beta} = 0, \quad (10)$$

with $\theta = 3\dot{a}/a$, a the scale factor of the FRW metric, a dot represents derivative wrt cosmological time, and h^α_λ is the metric of the 3-space orthogonal to v_μ . As shown in the pioneering work by Lifshitz [25], it is useful to expand the solutions of linear differential equations defined on a curved manifold in the corresponding spherical harmonics basis. In order to do so we will first split the equations in space and time parts. The vectors describing the electromagnetic field are already space vectors, since $E_\alpha V^\alpha = D_\alpha V^\alpha = B_\alpha V^\alpha = H_\alpha V^\alpha = 0$. From now on we shall work in conformal time, with the FRW metric given by

$$ds^2 = a(\eta)^2 [d\eta^2 - \gamma_{ij}(\vec{x}) dx^i dx^j],$$

where γ_{ij} is the metric of the hypersurface $\eta = \text{constant}$, and $i = 1, 2, 3$. In terms of 3-vectors, Eqns.(7)-(10) become

$$\nabla_i D^i = 0, \quad (11)$$

$$\nabla_i B^i = 0, \quad (12)$$

$$(D^i)' - \frac{a'}{3a} D^i + \eta^{jln} \nabla_l H_n = 0, \quad (13)$$

$$(B^i)' - \frac{a'}{3a} B^i - \eta^{jln} \nabla_l E_n = 0, \quad (14)$$

where the prime denotes derivative wrt conformal time.

Since we are dealing with vector quantities we will use a vector basis P^i defined by the following relations [26]

$$P^i = P^i(x^i),$$

$$\gamma^{ij} \nabla_i \nabla_j P^l = -m^2 P^l,$$

$$\gamma^{ij} \nabla_i P_j = 0$$

The eigenvalue m denotes the wave number of the corresponding vector eigenfunction of the Laplacian operator

on the spatially homogeneous hypersurfaces. The spectrum of eigenvalues depends on the 3-curvature ϵ , and is given by

$$m^2 = s^2 + 2, \quad 0 < s < \infty, \quad \epsilon = -1,$$

$$m = s, \quad 0 < s < \infty, \quad \epsilon = 0,$$

$$m^2 = s^2 - 2, \quad s = 2, 3, \dots, \quad \epsilon = 1.$$

The pseudovector basis P^i , convenient to work with the magnetic field, is defined by:

$$P^i = \eta^{ijl} \nabla_l P_j. \quad (15)$$

In terms of the basis P^i and of the associated basis $\tilde{P}^i$, the electric and magnetic fields can be expanded as follows:

$$E^i(\eta, \vec{x}) = \sum_{l, m, \sigma} E_{ml}^{(\sigma)}(\eta) P_{ml}^{(\sigma) i}(\vec{x}), \quad (16)$$

$$B^i(\eta, \vec{x}) = \sum_{l, m, \sigma} B_{ml}^{(\sigma)}(\eta) \tilde{P}_{ml}^{(\sigma) i}(\vec{x}). \quad (17)$$

These expressions are valid in the case $\epsilon = 1$. For $\epsilon = 0$ and -1 , the sum must be replaced by an integration $((2\pi)^{-3} \int d^3x)$. The index $l = 1, 2$ describes the two transverse degrees of freedom of the electric and magnetic fields. Since we are using an expansion in standing waves [38], the index σ takes the values "+" or "-". Consequently, the fields in the case $\epsilon = 0$ can be written as

$$E^i(\eta, \vec{x}) = \sum_{l, m} \left(E_{ml}^{(+)}(\eta) \cos(\vec{m} \cdot \vec{x}) + E_{ml}^{(-)}(\eta) \sin(\vec{m} \cdot \vec{x}) \right) e_l^i, \quad (18)$$

(where the e_l^i are modulus-one polarization vectors), and an analogous expression for B^i . In the more general case of $\epsilon = 1, -1$ the (considerably more involved) analytic expression for the vector base can be found in [27]. The result of substituting in Eqns.(11)-(14) the expansions given in Eqns.(16) and (17) is

$$B_{ml}^{(\sigma)'} + \frac{a'}{a} B_{ml}^{(\sigma)} - f^{-1}(\omega) D_{ml}^{(\sigma)} = 0, \quad (19)$$

$$D_{ml}^{(\sigma)'} + \frac{a'}{a} D_{ml}^{(\sigma)} + (m^2 + 2\epsilon) f(\omega) B_{ml}^{(\sigma)} = 0. \quad (20)$$

The system described by Eqns.(19) and (20) is not Hamiltonian when written in the variables D and B , but a Hamiltonian can be introduced in terms of a new set of variables defined by

$$p_{ml}^{(\sigma)}(\eta) \equiv a(\eta) D_{ml}^{(\sigma)}(\eta), \quad (21)$$

$$q_{ml}^{(\sigma)}(\eta) \equiv a(\eta) B_{ml}^{(\sigma)}(\eta). \quad (22)$$

In terms of these, Eqns.(19) and (20) can be written as

$$p_{ml}^{(\sigma)'} + (m^2 + 2\epsilon) f(\omega) q_{ml}^{(\sigma)} = 0, \quad (23)$$

$$q_{ml}^{(\sigma)'} - f(\omega)^{-1} p_{ml}^{(\sigma)} = 0. \quad (24)$$

Since the new variables constitute a pair of canonical variables, the dynamical system given by Eqns.(19) and (20) can now be described by the Hamiltonian

$$\mathcal{H}_{ml}^{(\sigma)}(p, q) = \frac{1}{2} b_2 p_{ml}^{(\sigma)2} + \frac{1}{2} b_1 q_{ml}^{(\sigma)2}, \quad (25)$$

where

$$b_1(\eta) = f(\omega)^{-1}, \quad b_2(\eta) = (m^2 + 2\epsilon) f(\omega). \quad (26)$$

The Hamiltonian for the mode characterized by (σ, m, l) given in Eqn.(25) is identical to that of a single harmonic oscillator problem with time-dependent coefficients. We shall review in the next section how to quantize this system.

II. QUANTIZATION

The expression for $\mathcal{H}$ obtained in the previous section appears in many different branches of physics, ranging from quantum optics to gravitational waves [28]. In order to quantize it, we shall follow a standard procedure of quantum optics. First we define creation and annihilation operators by the expressions

$$\hat{p}_{ml}^{(\sigma)} = -i\sqrt{\frac{\gamma}{2}} \left(\hat{a}_{ml}^{(\sigma)} - \hat{a}_{ml}^{(\sigma)\dagger} \right), \quad (27)$$

$$\hat{q}_{ml}^{(\sigma)} = \frac{\hat{a}_{ml}^{(\sigma)} + \hat{a}_{ml}^{(\sigma)\dagger}}{\sqrt{2\gamma}}, \quad (28)$$

where γ is a constant to be determined later. In terms of $\hat{a}$ and $\hat{a}^\dagger$, the Hamiltonian operator can be written as

$$\hat{\mathcal{H}}_{ml}^{(\sigma)} = \frac{1}{4} k(\eta) (2\hat{N}_{ml}^{(\sigma)} + 1) + \frac{1}{4} h(\eta) \hat{a}_{ml}^{(\sigma)2} + \frac{1}{4} h(\eta) \hat{a}_{ml}^{(\sigma)\dagger 2}, \quad (29)$$

where

$$k(\eta) = \gamma b_1(\eta) + \frac{b_2(\eta)}{\gamma}, \quad h(\eta) = \frac{b_2(\eta)}{\gamma} - \gamma b_1(\eta),$$

and $N_{ml}^{(\sigma)} = \hat{a}_{ml}^{(\sigma)\dagger} \hat{a}_{ml}^{(\sigma)}$ is the number operator. The first term in Eqn.(29) looks like the free Hamiltonian for the mode (σ, m, l) with a time-dependent function instead of the frequency. This part conserves the number of photons in each mode, hence the total number of photons

and the total energy. The second and third terms represent instead a time-dependent interaction which does not conserve the photon number.

The operators $\hat{a}_{ml}^\dagger$ and $\hat{a}_{ml}$ obey the commutation relations

$$[\hat{a}_{ml}^{(\sigma)}, \hat{a}_{m'l'}^{(\sigma')\dagger}] = \delta_{ll'} \delta_{mm'} \delta_{\sigma\sigma'}, \text{ for } \epsilon = 1, \quad (30)$$

$$[\hat{a}_{ml}^{(\sigma)}, \hat{a}_{m'l'}^{(\sigma')\dagger}] = \delta_{ll'} \delta_{\sigma\sigma'} \delta(m - m'), \text{ for } \epsilon = 0, -1, \quad (31)$$

where we have settled $\gamma = m$ in order to have vacuum as initial state. Since we shall work with a single mode (and in order to avoid clumsy expressions) the indices m, l and σ will be omitted from now onwards.

We will proceed to solve Schrödinger's equation for the Hamiltonian given in Eqn.(29) by writing the time evolution operator as a product of the rotation and the single-mode squeezed operators, along with a phase factor:

$$U(\eta, \eta_0) = e^{i\phi} S(\eta) R(\eta), \quad (32)$$

where $S(\eta)$ and $R(\eta)$ are defined as

$$S(\eta) = e^{A(\eta) \hat{a}^2 - A^*(\eta) \hat{a}^{\dagger 2}}, \quad (33)$$

$$R(\eta) = e^{i\Gamma(\eta) \hat{a}^\dagger \hat{a}}. \quad (34)$$

The function $\Gamma(\eta)$ is called the rotation angle, and $A(\eta)$ is usually defined as

$$A(\eta) = \frac{1}{2} r(\eta) e^{-2i\varphi(\eta)}, \quad (35)$$

where r is the squeeze factor (defined in the range $0 \leq r < \infty$) and φ the squeeze angle (defined in the range $-\pi/2 \leq \varphi < \pi/2$). The parameter r determines the strength of the squeezing while φ gives the distribution of the squeezing between conjugate variables. Schrödinger's equation with the Hamiltonian given in Eqn.(29) reduces, after a direct but somewhat long calculation, to the following system of first order coupled differential equations valid for each mode of the field:

$$ir' + (\varphi' + \Gamma') \sinh 2r = e^{2i\varphi} \frac{h}{2}, \quad (36)$$

$$-ir' + (\varphi' + \Gamma') \sinh 2r = e^{-2i\varphi} \frac{h}{2}, \quad (37)$$

$$(\varphi' + \Gamma') \cosh 2r - \varphi' = k, \quad (38)$$

$$\phi' = -\frac{\Gamma'}{2}. \quad (39)$$

Similar equations were obtained by Albrecht *et al*[29] in the case of the scalar degrees of freedom of metric perturbations in cosmology. Grishchuk [31] derived analogous

equations for the case of gravitational waves, and Matacz [32], for a scalar field. To solve the system, we shall follow the method presented in [32]. With the change of variables

$$\alpha(\eta) = e^{-i\Gamma(\eta)} \cosh r(\eta), \quad (40)$$

$$\beta(\eta) = -e^{-2i(\varphi(\eta) + \Gamma/2)} \sinh r(\eta), \quad (41)$$

Eqns.(36)-(39) can be written as

$$2\alpha' = -ih\beta - ik\alpha, \quad (42)$$

$$2\beta' = ik\beta + ih\alpha. \quad (43)$$

Introducing the function $\mu(\eta)$ defined by

$$\mu = \frac{\beta^* - \alpha^*}{\beta^* + \alpha^*}, \quad (44)$$

Eqns.(42) and (43) are equivalent to

$$2\mu' - i(k - h)\mu^2 + i(k + h) = 0. \quad (45)$$

Finally, setting

$$\mu = \frac{i}{\gamma b_1} \frac{g'}{g}, \quad (46)$$

we obtain the equation

$$g'' - \frac{b_1'}{b_1} g' + b_1 b_2 g = 0.$$

The problem has been reduced then to solving a single second-order differential equation for the function $g(\eta)$, which in terms of $f(\omega)$ reads (see Eqn(26))

$$g'' + \frac{1}{f} \frac{df}{d\omega} \omega' g' + (m^2 + 2\epsilon)g = 0. \quad (47)$$

We can see from this equation that there is no photon creation if the scalar field is constant, a fact that justifies the identification of ω with the laser “pump” used in experiments devised to observe squeezed states [28]. With the function g we can get the squeeze parameter r through Eqn.(46) as follows. From Eqns (40), (41), and (44) we get

$$\mu(\eta) = \frac{1 + e^{2i\varphi(\eta)} \tanh r(\eta)}{-1 + e^{2i\varphi(\eta)} \tanh r(\eta)}, \quad (48)$$

which can be inverted to get r , given by

$$\tanh^2 r(\eta) = \frac{1 + \mu(\eta) + \mu^*(\eta) + |\mu(\eta)|^2}{1 - \mu(\eta) - \mu^*(\eta) + |\mu(\eta)|^2}. \quad (49)$$

All the functions appearing in the evolution operator can be written in terms of μ (we refer the reader to [32] for details). A quantity of interest in the following is the mean number of photons per mode as a function of the conformal time, given by [35]

$$\langle N(\eta) \rangle = \sinh^2 r(\eta). \quad (50)$$

In the next sections we shall calculate $\langle N \rangle$ for two different models.

III. A SIMPLIFIED MODEL

Before going into the details of the model which takes into account the effect of matter creation on the metric, we shall study the production of photons in a simpler case, namely that in which the sole matter content is given by the scalar field ω . As discussed in [33], there are nonsingular solutions in the theory of WIST that describe a FRW geometry plus a Weyl scalar field. Nonsingular solutions are also present in string theory [34]. Let us briefly review these solutions. The EOM for gravitation plus scalar field written in conformal time are [33]

$$a'^2 + \epsilon a^2 + \frac{\lambda^2}{6}(\omega' a)^2 = 0, \quad (51)$$

$$\omega' = \sigma a^{-2},$$

where $\sigma = \text{constant}$, and λ^2 is the coupling constant of the scalar field to gravity. Notice that the equation of state of the scalar field is given by $\rho_\omega = p_\omega$, where

$$\rho_\omega = -\frac{\lambda^2}{2} \left(\frac{\omega'}{a} \right)^2.$$

From these equations we get

$$a'^2 = -\epsilon a^2 - \frac{b^2}{a^2}, \quad (52)$$

where we have defined $b^2 = \lambda^2 \sigma^2 / 6$. Eqn.(52) shows that only solutions with $\epsilon = -1$ are possible. Hence, from now on we shall restrict to the negative curvature case, for which Eqn.(52) can be easily integrated. Although this choice is at odds with observation, we emphasize that we are using this exact solution only as a toy model to study the effect of matter creation on the production of photons. The result of the integration for the scale factor is [39].

$$a(\eta) = \sqrt{|b|} \sqrt{\cosh(2\eta + \delta)}. \quad (53)$$

From Eqn.(51),

$$\omega(\eta) = \pm \frac{\sqrt{6}}{2\lambda} \arctan(e^{2\eta + \delta}) + \frac{\pi}{4} \frac{\text{sqr}t{6}}{\lambda}, \quad (54)$$

where δ is an integration constant. The plots for these functions are given in Fig.1. The scale factor displays a bounce, produced by the violation of the strong energy condition by the scalar field [37]. The number of photons created by the expansion in this model must be calculated numerically, since Eqn.(47) has no analytical solution for the scalar field given in Eqn.(54) with the coupling $f(\omega) = e^{-2\omega}$. The results are given by the dashed curve of Fig.(2), for a wavenumber typical of the size of the intergalactic scale (1 Mpc).




FIG. 1: Plot of the scale factor (full line) and ω as a function of the conformal time for $\lambda = |b| = 1$.




FIG. 2: Plot of the mean number of photons as a function of the conformal time for $m = 20$ for the case without matter (dashed line) and for the case with matter creation (full line, see next section), for $\lambda = 1$.

IV. A MODEL WITH MATTER CREATION

We shall consider in this section the production of photons in a background of the system composed of the scalar field plus matter and geometry, using a non-singular new solution that incorporates the effect of the creation of matter on the geometry. Friedmann's equation in conformal time for this case is given by

$$a'^2 - a^2 = -\frac{\lambda^2}{6}(\omega' a)^2 + \frac{a^4}{3}\rho_m, \quad (55)$$




FIG. 3: Plot of a and ω for $k_0 = 1/7$ and $a_0 = 0.93$, values chosen by imposing that the solution in Eqn.(59) enters the radiation era for $t \approx 10^{-8}$ seg.

while the second Einstein equation is

$$-3 \left(2 \frac{a''}{a^3} - \frac{a'^2}{a^4} - \frac{1}{a^2} \right) = \rho_m + 3\rho_\omega. \quad (56)$$

The conservation of the stress-energy tensor gives

$$\frac{d}{d\eta}(a^3(\rho_m + \rho_\omega)) + (p_m + p_\omega) \frac{da^3}{d\eta} = 0. \quad (57)$$

In the case of ultrarelativistic matter, this equation takes the form

$$(a^4 \rho_m)' + \frac{1}{a^2} (a^6 \rho_\omega)' = 0. \quad (58)$$

We would like to have a solution that describes creation of relativistic matter only around the bounce, and enters a radiation phase with a constant scalar field in a short time. Clearly, an asymmetry is to be expected both in the scale factor and in ω , since the evolution of this universe starts from the vacuum and enters a radiation-dominated epoch. An expression for a that fulfills these requirements is

$$a(\eta) = \beta \sqrt{\cosh(2\eta) + k_0 \sinh(2\eta) - 2k_0(\tanh(\eta) + 1)}, \quad (59)$$

with $\beta = a_0 \sqrt{\frac{1}{1-2k_0}}$ and $0 < k_0 \leq 1/7$. The expression for ω can be obtained using this scale factor in Eqns.(55) and (56). We shall not give the explicit expression, but the plots for $a(\eta)$ and $\omega(\eta)$ (see Fig.3). Notice that the plot shows the announced asymmetry. The evolution of the Hubble parameter for the vacuum case (studied in the previous section) and for the case with matter creation is showed in Fig.4. Let us emphasize that since the




FIG. 4: Plot of H for the vacuum case (full line) and for the solution given by Eqn.(59) as a function of the conformal time for $\lambda = 1$ in $c = 1$ units.

scalar field tends rapidly to a constant value, the production of matter (controlled by ω' , see Eqn.(58)) stops soon, and the model enters a radiation phase *without the need of a potential*. In this sense, this solution describes a *hot bounce*, as opposed to cold bouncing solutions, which do not enter the radiation era unless they are heated up [36]. Another nice feature of this solution is that the dilaton goes automatically to a constant value for $\eta \rightarrow \infty$, in such a way that the solution could be taken as the leading order of a perturbative development (as is the case in string theory). Again, no potential was needed in order to display this feature. The mean number of photons per mode can be calculated as in the previous cases, after numerical integration of Eqn.(47). The full line plot in Fig.2 shows that the production of photons is increased in the case of the model with matter creation. The number of photons is directly related to the fraction of electromagnetic energy stored per mode, relative to the background radiation energy ρ_γ through the expression

$$r(\nu) \approx \frac{\nu^4}{\rho_\gamma} \langle N \rangle_\nu,$$

where $\nu = \sqrt{m^2 - 2/a}$. Consequently, the increment in the number of photons may be of importance in the problem of the generation of magnetic field seeds.

V. DISCUSSION

Using a covariant description for the electromagnetic field coupled to a scalar in a curved background, we obtained the Hamiltonian for each mode of the field. With the aid of the formalism of the squeezed states, we have

calculated the number of photons in a non-singular universe using first an already known solution [33] with no matter present, and afterwards a new solution which takes into account the effect of the creation of ultrarelativistic matter on the evolution of the scale factor and of the scalar field. This new solution presents several interesting features, namely the transition of an expanding phase to a radiation era, and the constancy of the scalar field a short time after the bounce. The results for the mean photon number per mode show that the production is increased in the case of matter production. This

increment may be relevant for the creation of seeds of the magnetic field. We hope to discuss in detail this issue in a forthcoming publication.

Acknowledgements

SEPB acknowledges UERJ for financial support. JS is supported by CNPq.

-
- [1] E. Parker, Ap. J. **162**, 665 (1970). See also Y. B. Zel'dovich, A. A. Ruzmaikin, and D. D. Sokoloff, *Magnetic Fields in Astrophysics*, Mc Graw-Hill, New York (1980).
 - [2] See however R. Kulsrud, S. C. Cowley, A. V. Gruzinov and R. N. Sudan, Phys. Rep. **283**, 213(1997), and L. M. Widrow, Rev. Mod. Phys. **74**, 775 (2003).
 - [3] L. Biermann, Z. Naturf. 5A, 65 (1950).
 - [4] T.E. Clarke, P.P. Kronberg and H. Böhringer, Astrophys. J. **547**, L111 (2001).
 - [5] H. Böhringer, Rev. Mod. Astron. **8**, 295 (1995).
 - [6] S. A. Colgate and H. Li, astro-ph/0001418.
 - [7] D. Grasso and H. Rubinstein, Phys. Rept. **348**, 163 (2001).
 - [8] M. Turner and L. Widrow, Phys. Rev. D **37**, 2743 (1988).
 - [9] For a review on different mechanisms of seed generation, see *Generation of magnetic fields in cosmology*, A. Dolgov, in Gurzadyan, V.G. (ed.) et al.: *From integrable models to gauge theories*, 143-154, hep-ph/0110293.
 - [10] *Gauge invariant wave equations in curved space-times and primordial magnetic fields*, G. Lambiase and A. Prasanna, gr-qc/0407071.
 - [11] M. Novello and J. Salim, Phys. Rev. D **20**, 377 (1978).
 - [12] A. Dolgov, Phys. Rev. D **48**, 2499 (1993).
 - [13] E. Calzetta, A. Kandus and F. Mazzitelli, Phys. Rev. D **57**, 7139 (1998).
 - [14] B. Ratra, Astrophys. J. **391**, L1 (1992).
 - [15] *Nearly minimal magnetogenesis*, T. Prokopec and Ewald Puchwein, astro-ph/0403335.
 - [16] Lemoine D., Lemoine M., Phys. Rev. D **52**, (1995), 1955, Gasperini M., Giovannini M. and Veneziano G. Phys. Rev. Lett. **V 75**, (1995), 3796.
 - [17] J. Lidsey, D. Wands, E. Copeland, Phys. Rept. **337**, 343 (2000), hep-th/9909061.
 - [18] J. M. Salim and S. Sautu, Class. Quant. Gravity, **13** (1996) 353-360, J. M. Salim, S. Sautu and Martins R., Class. Quant. Gravity, **15** (1998) 1521.
 - [19] P. Dirac, Nature **139**, 323 (1937).
 - [20] See *Magnetogenesis, variation of gauge couplings, and inflation*, M. Giovannini, Proceedings of the Chalonge School on Astrofundamental Physics, N.G. Sanchez and Y.M. Pariiski, eds. Kluwer (2002), astro-ph/0212346.
 - [21] K. Bamba and J. Yokoyama, Phys. Rev. D **69**, 043507 (2004), astro-ph/0310824.
 - [22] *Large-scale magnetic fields from dilaton inflation in non-commutative spacetime*, K. Bamba and J. Yokoyama, hep-ph/0409237. See also *Generation of Cosmological seed magnetic fields from inflation with cutoff*, A. Ashoorioon and R. Mann, gr-qc/0410053.
 - [23] See for instance D. Bailin and A. Love, Rept. Prog. Phys. **50**, 1087 (1987).
 - [24] See for instance T. Damour and A. M. Polyakov, Nucl. Phys. **B423**, 532 (1994), hep-th/9401069.
 - [25] Lifshitz E. M. and Khalatnikov I. M., Adv. Phys. **12**, (1963), 185.
 - [26] M. Novello, J. M. Salim, M.C. Motta da Silva, S. E. Joras and R. Klippert, Phys. Rev. D **52**, (1995), 730.
 - [27] E. Kalnins and W. Miller, Jr., J. Math. Phys **32**, 698 (1991).
 - [28] B. Schumaker, Phys. Rep. **135**, 317 (1986).
 - [29] A. Albrecht, P. Ferreira, M. Joyce, T. Prokopec, Phys. Rev. D **50**, 4807 (1994), astro-ph/9303001.
 - [30] L. Grischuk and Y. Sidorov, Phys. Rev. D **42**, 3413 (1990).
 - [31] L. Grischuk, Phys. Rev. D **45**, 4717 (1992).
 - [32] Mataraz A. L., Phys. Rev. D **49**, (1994), 788.
 - [33] M. Novello, L. Oliveira, J.M. Salim and E. Elbaz, Int. J. of Mod. Phys. D **1**, 641 (1993).
 - [34] See for instance *Old ideas and new twists in string cosmology*, M. Giovannini, hep-th/0409251.
 - [35] M. Giovannini, *Primordial Magnetic Fields*, hep-ph/0208152.
 - [36] M. Giovannini, Class. Quantum Grav. **21**, 4209 (2004).
 - [37] C. Molina-Paris and M. Visser Phys. Lett. B **455**, 90 (1999).
 - [38] Notice that the expansion in standing waves is completely equivalent to that of travelling waves (see for instance [30]).
 - [39] The expression for the scale factor in terms of cosmological time was given in [33], although not in a closed form